

Quaternary Prevention: Need of the Hour

ABSTRACT

Prevention is primarily categorized as Primordial, Primary, Secondary and Tertiary. Now the concept of “Quaternary Prevention” is also introduced. This editorial article discusses need of Quaternary prevention in current scenario of clinical practice.

Keywords: Prevention, levels, quaternary

It is always said “prevention is better than cure.” This principle is undisputed. Preventive measures can be applied at any stage along the natural history of the disease with the goal of preventing further progression of the condition. Levels of the prevention are mainly categorized as primordial, primary, secondary, and tertiary prevention. Over the years, the concept of prevention has undergone significant changes.

Primordial prevention consists of actions to minimize future hazards to health and hence inhibits the establishment of factors which are known to increase the risk of disease. It addresses broad health determinants rather than preventing personal exposure to risk factors, which is the goal of primary prevention. Primary prevention seeks to prevent the onset of specific diseases via risk reduction by altering behaviors or exposures that can lead to disease or by enhancing resistance to the effects of exposure to a disease agent. Secondary prevention includes procedures that detect and treat preclinical pathological changes and thereby control disease progression. Screening procedures are often the first step, leading to early interventions that are more cost effective than intervening once symptoms appear. Once the disease has developed and has been treated in its acute clinical phase, tertiary prevention seeks to soften the impact caused by the disease on the patient’s function, longevity, and quality-of-life. For reversible conditions, tertiary prevention will reduce the population prevalence, whereas for incurable conditions it may increase the prevalence if it prolongs survival. While where the condition is not reversible, tertiary prevention focuses on rehabilitation, assisting the patient to accommodate to his disability.^[1]

Quaternary prevention a “higher” level of prevention has also been suggested first by Jamouille and Ronald. Quaternary prevention is defined as “action taken to identify patient at risk of over-medicalization, to protect him from new medical invasion, and to suggest him interventions ethically acceptable.” The concept of quaternary prevention makes it easier to “identify patient at risk of over-medicalization.”^[2] Health activities not only generally produce benefits, but also

harm. That is to say, although medical intervention is mainly favorable, there is a dynamic balance that requires continuous assessment of the clinical situation as naturally only those health activities that achieve more benefit than harm at the end are justified.^[3] Quaternary prevention is also essential in the phenomenon called disease mongering, which could be translated as commercialization of disease. Encouraging this sort of prevention and halting the consequences of disease mongering requires the development of all the institutional potential for prevention, as well as all the personal willingness for restraint; it involves separating us from the unnecessary auspices of industry, being critical of our work, not being maleficent, respecting the principle of justice as managers of the limited public resources and making ourselves feel responsible for the social cost resulting from medical decisions.^[4]

To conclude, quaternary prevention has become an essential component of prevention concept and it should be applied as a prevention tool in current scenario in many disease conditions pertaining to classic “primum non nocere” meaning “first, do no harm.”

Harshal T. Pandve

Department of Community Medicine, Smt. Kashibai Navale Medical College, Narhe, Pune - 411 041, Maharashtra, India

Address for correspondence: Dr. Harshal T. Pandve,
Department of Community Medicine, Smt. Kashibai Navale
Medical College, Narhe, Pune - 411 041, Maharashtra, India.
E-mail: dr_harshalpandve@yahoo.co.in


References

1. Available from: <http://www.phprimer.afmc.ca/Part1-TheoryThinkingAboutHealth/Chapter4BasicConceptsInPreventionSurveillanceAndHealthPromotion/Thestagesofprevention>. [Last accessed on 2014 Jun 27].
2. Jamouille M, Roland M. Quaternary prevention. Paper presented at the Hong-Kong Meeting of the Wonca Classification Committee; June, 1995.

Pandve: Quaternary prevention: Need of the hour

3. Gervas J. Quaternary prevention in the elderly. *Rev Esp Geriatr Gerontol* 2012;47:266-9.
4. Martínez González C, Riaño Galán I, Sánchez Jacob M, González de Dios J. Quaternary prevention: Containment as an ethical necessity. *An Pediatr (Barc)* 2014;pii: S1695-403300283-5.

Access this article online	
Quick Response Code:	
	Website: www.jfmfc.com



Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

- 1) **First Page File:**
Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.
- 2) **Article File:**
The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.
- 3) **Images:**
Submit good quality color images. Each image should be less than 4096 kb (4 MB) in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.
- 4) **Legends:**
Legends for the figures/images should be included at the end of the article file.